

**Phase I, 0-2 weeks.** Pain and edema control, start flexibility and strengthening exercises.

PRECAUTIONS	<ul style="list-style-type: none"> <li>- Do not put a pillow under the operated knee for comfort when elevating LE, i.e. maintain full knee extension</li> <li>- You may bear full weight with the brace locked in extension</li> <li>- Use crutches until gait and balance have normalized</li> <li>- Avoid heat application</li> <li>- Do not bend the knee beyond 90°</li> <li>- Avoid lateralization of patella</li> <li>- The brace remains locked in extension for ambulation and sleep for 4 weeks</li> </ul>
Emphasize	<ul style="list-style-type: none"> <li>- Patellar mobility (no lateral)</li> <li>- Full PROM knee extension</li> <li>- Improving quadriceps contraction</li> <li>- Controlling pain and swelling</li> <li>- Compliance with HEP and precautions</li> </ul>
Assessment	<ul style="list-style-type: none"> <li>- LEFS, IKDC, SANE, ACL RSI, NPRS</li> <li>- Swelling</li> <li>- Girth measurements</li> <li>- Neurovascular assessment</li> <li>- Wound status</li> <li>- Patellar mobility</li> <li>- Quality of quadriceps contraction</li> <li>- LE AROM and PROM</li> <li>- Straight leg raise (SLR) in supine</li> </ul>
Treatment Recommendations	<ul style="list-style-type: none"> <li>- Passive knee extension with towel under heel</li> <li>- Quadriceps re-education: quadriceps sets with towel under knee with neuromuscular electric stimulation (NMES) or biofeedback</li> <li>- Patellar mobilization (no lateral)</li> <li>- AROM knee flexion to 90°, AROM knee extension to 0°</li> <li>- PROM knee flexion no more than 90°</li> <li>- SLR in all planes (With brace locked at 0° in supine)</li> <li>- Hip progressive resistive exercises</li> <li>- Calf strengthening (Unilateral elastic band)</li> <li>- Upper extremity ergometry, as tolerated</li> <li>- Edema management, e.g. cryotherapy (no submersion), elevation, gentle edema mobilization avoiding incision</li> <li>- Progressive home exercise program</li> </ul>
Criteria for Advancement	<ul style="list-style-type: none"> <li>- Ability to SLR without quadriceps lag or pain</li> <li>- Knee ROM 0°-90°</li> <li>- Pain and swelling controlled</li> </ul>

**Phase II, 2-6 weeks.** Progressive strengthening while protecting the repair.

PRECAUTIONS	<ul style="list-style-type: none"> <li>- Do not put a pillow under the operated knee for comfort when elevating LE, i.e. maintain full knee extension</li> <li>- You may bear full weight with the brace locked in extension</li> <li>- Use crutches until gait and balance have normalized</li> <li>- Avoid heat application</li> <li>- Avoid lateralization of patella</li> <li>- The brace remains locked in extension for ambulation and sleep for 4 weeks</li> </ul>
Emphasize	<ul style="list-style-type: none"> <li>- Patellar mobility (no lateral)</li> <li>- Full PROM knee extension</li> <li>- Improving quadriceps contraction</li> <li>- Controlling pain and swelling</li> <li>- Compliance with HEP and precautions</li> </ul>
Assessment	<ul style="list-style-type: none"> <li>- LEFS, IKDC, SANE, ACL RSI, NPRS</li> <li>- Swelling</li> <li>- Girth measurements</li> <li>- Gait and Neurovascular assessment</li> <li>- Wound status</li> <li>- Patellar mobility</li> <li>- Quality of quadriceps contraction</li> <li>- LE AROM and PROM</li> <li>- Straight leg raise (SLR) in supine</li> <li>- Single leg stance, when appropriate (around 6 weeks)</li> </ul>
Treatment Recommendations	<ul style="list-style-type: none"> <li>- Passive knee extension with towel under heel</li> <li>- Quadriceps re-education: quadriceps sets with towel under knee with neuromuscular electric stimulation (NMES) or biofeedback</li> <li>- Patellar mobilization</li> <li>- AROM knee flexion to 90°</li> <li>- Knee PROM to 90°</li> <li>- Progression from seated to standing (wall slides) to bike ROM</li> <li>- Straight leg raises (SLR) PRE's in all planes             <ul style="list-style-type: none"> <li>• With brace locked at 0° in supine until no extension lag</li> </ul> </li> <li>- Leg press bilaterally in 80°-5° arc after 4 weeks             <ul style="list-style-type: none"> <li>• Progression from bilaterally, to 2 up/1 down, to unilateral</li> </ul> </li> <li>- Functional strengthening             <ul style="list-style-type: none"> <li>• Mini squats up to 0°-60° after 4 weeks</li> <li>• Forward step-up progression starting with 2"-4" (around 5 weeks)</li> </ul> </li> <li>- Terminal knee extension in weight bearing</li> <li>- Consider blood flow restriction (BFR) program</li> <li>- Hip-gluteal progressive resistive exercises             <ul style="list-style-type: none"> <li>• May introduce Romanian Dead Lift toward end of phase</li> </ul> </li> <li>- Hamstring strengthening</li> <li>- Calf strengthening (Progression to unilateral calf raises)</li> <li>- Flexibility exercises</li> <li>- Proprioception board/balance system             <ul style="list-style-type: none"> <li>• Progression from bilateral to unilateral weight bearing</li> </ul> </li> </ul>

MPFL RECONSTRUCTION REHABILITATION PROTOCOL

	<ul style="list-style-type: none"><li>- Stationary bicycle<ul style="list-style-type: none"><li>• Short crank for ROM and/or cycling (&lt;90° knee flexion)</li></ul></li><li>- Upper extremity ergometry, as tolerated</li><li>- Gait training WBAT around 5-6 weeks</li><li>- Edema management, e.g. cryotherapy (no submersion until incision adequately healed), elevation, gentle edema mobilization avoiding incision</li><li>- Progressive home exercise program</li><li>- Patient education regarding monitoring of response to increase in activity level and weightbearing</li></ul>
Criteria for Advancement	<ul style="list-style-type: none"><li>- Knee ROM 0°-90°</li><li>- Good patellar mobility</li><li>- Minimal swelling</li><li>- Full weight bearing without pain</li></ul>

**Phase III, 7-12 weeks.** Regain functional movement and strength.

PRECAUTIONS	<ul style="list-style-type: none"> <li>- Discontinue brace when stability, balance, and control achieved</li> <li>- Avoid pain with exercises, standing, walking and other activities</li> <li>- Monitor tolerance to load, frequency, intensity and duration</li> <li>- Avoid too much too soon</li> </ul>
Emphasize	<ul style="list-style-type: none"> <li>- Address impairments</li> <li>- Functional movement and strength</li> <li>- Activity level to match response and ability</li> </ul>
Assessment	<ul style="list-style-type: none"> <li>- LEFS, IKDC, SANE, ACL RSI, NPRS</li> <li>- Swelling</li> <li>- Girth measurements</li> <li>- Gait and Neurovascular assessment</li> <li>- Wound status</li> <li>- Patellar mobility</li> <li>- Quality of quadriceps contraction</li> <li>- LE AROM and PROM</li> <li>- Straight leg raise (SLR) in supine</li> <li>- Functional assessment, e.g. single leg stance, step ups/downs, squat, gait</li> <li>- Balance testing, e.g. Star Excursion Test, Biodex Balance System<sup>TM</sup></li> <li>- Quadriceps isometrics testing with dynamometer at 60° at 12 weeks</li> </ul>
Treatment Recommendations	<ul style="list-style-type: none"> <li>- Patellar mobilization</li> <li>- ROM 0-110° (8 wks), 120° (10 wks), then full</li> <li>- SLR PRE's in all planes</li> <li>- Leg press eccentrically</li> <li>*Functional strengthening             <ul style="list-style-type: none"> <li>• Progress squats to 0°-90°, initiating movement with hips</li> <li>• Continue forward step-up progression</li> <li>• Initiate step-down progression starting with 2"-4"</li> <li>• Lateral step-ups, crossovers</li> <li>• Lunges</li> </ul> </li> <li>- Continue foundational hip-gluteal progressive resistive exercises</li> <li>- Continue hamstring and calf strengthening</li> <li>- Flexibility exercises and foam rolling</li> <li>- Core and UE strengthening</li> <li>- Consider BFR program</li> <li>* Proprioception training             <ul style="list-style-type: none"> <li>• Continue foundational exercises</li> <li>• Progress to perturbation training</li> </ul> </li> <li>* Cardiovascular conditioning             <ul style="list-style-type: none"> <li>• Stationary bicycle</li> <li>• Elliptical when able to perform 6" step-up with good form</li> </ul> </li> <li>- Gait training WBAT</li> <li>- Cryotherapy</li> </ul>

Dr. Tyler Kent  
Tylerkentmd.com

MPFL RECONSTRUCTION REHABILITATION PROTOCOL

	<ul style="list-style-type: none"><li>• Ice with passive knee extension with towel under heel as needed to maintain ROM</li><li>- Progressive home exercise program</li><li>- Patient education regarding monitoring of response to increase in activity level</li></ul>
Criteria for Advancement	<ul style="list-style-type: none"><li>- Ability to perform 8" step-down with good control and alignment without pain</li><li>- Full symmetrical knee ROM</li><li>- Symmetrical squat to parallel</li><li>- Single leg bridge holding for 30 seconds</li><li>- Balance testing and quadriceps isometrics 70% of contralateral lower extremity</li></ul>

**Phase IV, 13-19 weeks.** Begin job- or sport-specific training.

PRECAUTIONS	<ul style="list-style-type: none"> <li>- Initiate return to running/sport only when cleared by physician</li> <li>- Avoid pain with exercises and functional training</li> <li>- Monitor tolerance to load, frequency, intensity and duration</li> <li>- Avoid too much too soon</li> </ul>
Emphasize	<ul style="list-style-type: none"> <li>- Address impairments</li> <li>- Return to normal functional activities</li> </ul>
Assessment	<ul style="list-style-type: none"> <li>- LEFS, IKDC, SANE, ACL RSI, NPRS</li> <li>- Swelling</li> <li>- Girth measurements</li> <li>- LE AROM and PROM</li> <li>- Functional assessment, e.g. single leg stance, step ups/downs, squat, gait</li> <li>- Balance testing, e.g. Star Excursion Test, Biodex Balance System™</li> <li>- Quadriceps isometrics or isokinetic testing</li> <li>- QMA – Quality of Movement Testing</li> </ul>
Treatment Recommendations	<ul style="list-style-type: none"> <li>- Progress leg press eccentrically</li> <li>* Functional strengthening             <ul style="list-style-type: none"> <li>• Progress squats to 0°-90°, initiating movement with hips</li> <li>• Progress to single leg squats</li> <li>• Forward step-up and step-down progression</li> <li>• Progress lateral step-ups, crossovers</li> <li>• Progress lunges</li> </ul> </li> <li>- Initiate running progression</li> <li>- Initiate plyometric progression</li> <li>- Continue foundational hip-gluteal progressive resistive exercises</li> <li>- Continue hamstring and calf strengthening</li> <li>- Flexibility exercises and foam rolling</li> <li>- Core and UE strengthening</li> <li>- Consider BFR program</li> <li>* Progress proprioception training             <ul style="list-style-type: none"> <li>• Continue foundational exercises</li> <li>• Incorporate agility and controlled sports-specific movements</li> </ul> </li> <li>* Progress cardiovascular conditioning             <ul style="list-style-type: none"> <li>• Stationary bicycle</li> <li>• Elliptical</li> </ul> </li> <li>- Cryotherapy and/or compression therapy</li> <li>- Progressive home exercise program</li> <li>- Patient education regarding monitoring of response to increase in activity level</li> </ul>
Criteria for Advancement	<ul style="list-style-type: none"> <li>- No swelling</li> <li>- Normal neurovascular assessment</li> <li>- Normal scar and patellar mobility</li> <li>- Normal quadriceps contraction</li> <li>- Full LE ROM, flexibility and strength</li> <li>- Quantitative assessments = 85% of contralateral lower extremity</li> </ul>

**Phase V, weeks 20 - discharge.** Return to full activity without impairment or restrictions.

PRECAUTIONS	<ul style="list-style-type: none"> <li>- Note importance of gradual return to participation with load and volume monitoring under guidance of physical therapist, MD, athletic trainer and coach</li> <li>- Avoid premature or too rapid full return to sport</li> </ul>
Emphasize	<ul style="list-style-type: none"> <li>- Return to participation</li> <li>- Collaboration with Sports Performance experts</li> </ul>
Assessment	<ul style="list-style-type: none"> <li>- LEFS, IKDC, SANE, ACL RSI, NPRS</li> <li>- Swelling</li> <li>- LE AROM and PROM</li> <li>- Balance testing, e.g. Star Excursion Test, Biodex Balance System™</li> <li>- Quadriceps isometrics or isokinetic testing</li> <li>- Functional and Return to Sport tests, e.g. hop testing, QMA – Quality of Movement Testing</li> </ul>
Treatment Recommendations	<ul style="list-style-type: none"> <li>- Gradually increase volume and load to mimic load necessary for return to activity</li> <li>- Progress movement patterns specific to patient's desired sport or activity</li> <li>- Progression of agility work</li> <li>- Increase cardiovascular load to match that of desired activity</li> <li>- Collaborate with ATC, performance coach/strength and conditioning coach, skills coach and/or personal trainer to monitor load and volume as return to participation</li> <li>- Consult with referring MD on timing return to sport including any recommended limitations</li> </ul>
Criteria for Return to Sport	<ul style="list-style-type: none"> <li>- Quantitative assessments = 90% of contralateral lower extremity</li> <li>- Movement patterns, functional strength, flexibility, motion, endurance, power, deceleration and accuracy to meet demands of sport</li> </ul>